



References for Probiotic Bulletin, Issue 26

Pages 1-3: Probiotics, stress and upper respiratory tract infections

1. McCormick A, Fleming D and Charlton J (1995) Royal College of General Practitioners, Office of Population Censuses and Surveys, and Department of Health. *Morbidity statistics from General Practice. Fourth National Survey 1991-1992*. London: HMSO.
2. Easton G, Saxena A (2010) Antibiotic prescribing for upper respiratory tract infections in children: how can we improve? *London J Primary Care* 3:37-41.
3. Fahey T, Stocks N and Thomas T (1998) Systematic review of the treatment of upper respiratory tract infection. *Arch Dis Child* 79, 225-230.
4. Ashworth M, Latinovic R, Charlton J et al (2004) Why has antibiotic prescribing for respiratory illness declined in primary care? A longitudinal study using the General Practice Research Database. *J Public Health* 26(3):268-274.
5. Gulliford MC, Dregan A, Moore MV et al (2014) Continued high rates of antibiotic prescribing to adults with respiratory tract infection: survey of 568 UK general practices. *BMJ Open* 4:e006245.
6. Wallinga D, Rayner G, Lang T (2015) Antimicrobial resistance and biological governance: explanations for policy failure. *Public Health* 129:1314-1325.
7. Office for National Statistics (2014) Sick leave in the labour market: February 2014. Analysis describing sickness absence rates of employees in the labour market. http://webarchive.nationalarchives.gov.uk/20160105160709/http://ons.gov.uk/ons/dcp171776_353899.pdf
8. Shida K, Sato T, Iizuka R et al (2015) Daily intake of fermented milk with *Lactobacillus casei* strain Shirota reduces the incidence and duration of upper respiratory tract infections in healthy middle-aged office workers. *European Journal of Nutrition* [Epub ahead of print].
9. Cohen S, Doyle WJ, Skoner DP (1999) Psychological stress, cytokine production, and severity of upper respiratory illness. *Psychosom Med* 61:175-180.
10. Mayer EA, Tillisch K, Gupta A (2015) Gut/brain axis and the microbiota. *The Journal of Clinical Investigation* 125:926-938.
11. Kato-Kataoka K, Nishida K, Takada M et al (2015) Fermented milk containing *Lactobacillus casei* Shirota prevents the onset of physical symptoms in medical students under academic examination stress. *Beneficial Microbes* 7(2):153-6.
12. Benton D, Williams C and Brown A (2007) Impact of consuming a milk drink containing a probiotic on mood and cognition. *Eur J Clinical Nutr* 61:355-361.
13. Rao AV, Bsted AC, Beaulne TM, et al (2009). A randomized, double-blind, placebo-controlled pilot study of a probiotic in emotional symptoms of chronic fatigue syndrome. *Gut Pathogens* 1:6.
14. Gleeson M, Bishop NC, Oliveira M, Tauler P (2011) Daily probiotic's (*Lactobacillus casei* Shirota) reduction of infection incidence in athletes. *Int J Sports Nutr Exerc Metab* 21:55-64.
15. Shida K and Nomoto K (2013) Probiotics as efficient immunopotentiators: Translational role in cancer prevention *Indian J Med Res* 139:808-819.
16. Gleeson M (2016) Immunological aspects of sports nutrition. *Immunol Cell Biol* 94:117-123.

17. [Fujita R, Iimuro S, Shinozaki T, et al \(2013\)](#) Decreased duration of acute upper respiratory tract infections with daily intake of fermented milk: a multicenter, double-blinded, randomized comparative study in users of day care facilities for the elderly population. *Am J Infect Control* 41(12):1231-5.
18. van Puyenbroeck K, Hens N, Coenen S, et al (2012) Efficacy of daily intake of *Lactobacillus casei* Shirota on respiratory symptoms and influenza vaccination response: a randomized, double-blind, placebo-controlled trial in healthy elderly nursing home residents. *Am J Clin Nutr* 95:1165-1171.
19. King S, Glanville J, Sanders ME et al (2014) Effectiveness of probiotics on the duration of illness in healthy children and adults who develop common acute respiratory infectious conditions: a systematic review and meta-analysis. *Br J Nutr* 112:41-54.
20. Hao Q, Dong BR, Wu T (2015) Probiotics for preventing acute upper respiratory tract infections. *Cochrane Database Syst Rev* 2:CD006895.
21. Lenoir-Wijnkoop I, Gerlier L, Bresson J-L, et al (2015) Public health and budget impact of probiotics on common respiratory tract infections: a modelling study. *PLoS ONE* 10(4):e0122765.
22. Bhangar S, Adams RI, Pasut W, Huffman JA, Arens EA, Taylor JW, Bruns TD, Nazaroff WW. 2015. Chamber bioaerosol study: human emissions of size-resolved fluorescent biological aerosol particles. *Indoor Air* 26:193-206.
23. You R, Cui W, Chen C, Zhao B. 2013. Measuring the short-term emission rates of particles in the "personal cloud" with different clothes and activity intensities in a sealed chamber. *Aerosol and Air Quality Research* 13:911–921.
24. Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, et al. Disordered microbial communities in asthmatic airways. *PLoS One*. 2010; 5: e8578.
25. Dickson RP, Erb-Downward JH, Huffnagle GB (2013) The role of the bacterial microbiome in lung disease. *Expert Rev Respir Med* 7:245-257.
26. Cystic Fibrosis Trust (2016) Breathe – Life Unlimited: We're fighting for every Breath. <https://www.cysticfibrosis.org.uk/>

Page 4: Probiotic potential for infection control in care homes

Bian L, Nagata S, Asahara T *et al* (2011) Effects of the continuous intake of *Lactobacillus casei* strain Shirota-fermented milk on risk management of long-term inpatients at health service facilities for the elderly. *Int J Probiotics Prebiotics* 6(2):123-132.

Gerba CP, Rose JB, Haas CN (1996) Sensitive populations: who is at the greatest risk? *Int J Food Microbiol* 30:113-123.

Nagata S, Asahara T, Ohta T *et al* (2011) Effect of the continuous intake of probiotic-fermented milk containing *Lactobacillus casei* strain Shirota on fever in a mass outbreak of norovirus gastroenteritis and the faecal microflora in a health service facility for the aged. *Br J Nutr* 106(4):549-56.

Pirker A, Stockenhuber A, Remely M *et al* (2013) Effects of antibiotic therapy on the gastrointestinal microbiota and the influence of *Lactobacillus casei*. *Food Agric Immunol* **24(3)**:315-330.

Van den Nieuwboer M, Klomp-Hogeterp A, Verdoorn S *et al* (2015) Improving the bowel habits of elderly residents in a nursing home using probiotic fermented milk. *Beneficial Microbes* 6(4):397-403.

Pages 6 – 7: The influence of the gut microbiota on weight gain and eating disorders

Armougom F, Henry M, Vialettes B et al (2009). Monitoring bacterial community of human gut microbiota reveals an increase in *Lactobacillus* in obese patients and methanogens in anorexic patients. *PLoS One* 4: e7125.

Gaci N, Borrel G, Tottey W et al (2014) Archaea and the human gut: new beginning of an old story. *World J Gastroenterol* 20(43):16062-78.

Goodrich JK, Waters JL, Poole AC et al (2014). Human genetics shape the gut microbiome. *Cell* 159:789–799.

Gouba N, Raoult D, Drancourt M (2014) Gut microeukaryotes during anorexia nervosa: a case report. *BMC Res Notes* 7:33.

JCPFMH (Joint Commissioning Panel For Mental Health) (www.jcpmh.info/wp-content/uploads/10keymsgs-eatingdisorders.pdf)

Mbakwa CA, Penders J, Savelkoul PH et al (2015) Gut colonization with *Methanobrevibacter smithii* is associated with childhood weight development. *Obesity (Silver Spring)* 23(12): 2508-16.

Morita C, Tsuji H, Hata T et al (2015) Gut dysbiosis in patients with anorexia nervosa. *PLoS ONE* 10(12):e0145274.

NICE Guidelines, 2004 page 7. "Eating Disorders Core Interventions in the Treatment and Management of Anorexia Nervosa, Bulimia Nervosa, and Related Eating Disorders". *National Collaborating Centre for Mental Health*.

Turnbaugh PJ, Hamady M, Yatsunencko T et al (2009a) A core gut microbiome in obese and lean twins. *Nature* 457(7228):480-4.

Turnbaugh PJ, Gordon JI. (2009b) The core gut microbiome, energy balance and obesity. *J Physiol.* Sep 1;587(Pt 17):4153-8. Sep 1;587(Pt 17):4153-8.

Wu GD, Chen J, Hoffmann C et al (2011) Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334:105-108.

For further information visit our website: www.yakult.co.uk/hcp
Contact the science team on science@yakult.co.uk or 0208 8427 600.