

YOUR GUIDE TO Fibre

Dietary fibres are carbohydrates that are not digested or absorbed in the small intestine and therefore reach the large intestine, where they can be broken down by bacteria.



DEFINITION

Dietary fibres, as defined by Codex¹ and the European Commission², are carbohydrate polymers with three or more monomeric units, which are neither digested nor absorbed in the small intestine and belong to the following categories:

- edible carbohydrate polymers naturally occurring in the food as consumed;
- edible carbohydrate polymers which have been obtained from food raw material by physical, enzymatic or chemical means and which have a beneficial physiological effect demonstrated by generally accepted scientific evidence;
- edible synthetic carbohydrate polymers and which have a beneficial physiological effect demonstrated by generally accepted scientific evidence.

DIETARY RECOMMENDATION

In the UK and Ireland, the fibre intake of adults is far below what is recommended, with an average daily intake of only 19g.^{3,4}

Table 1. UK and Ireland Daily Fibre Recommendations⁵⁻⁷

Age Group	UK	Ireland
2-4 years old	15g	Age +5g
5-10 years old	20g	Age +5g
11-14 years old	25g	Age +5g
15-18 years old	30g	Age +5g
19+ years old	30g	≥25g

FIBRE RICH FOODS

Dietary fibre is naturally present in many plant-based foods.

Here are some examples and the amount of fibre they provide (per 100g):

 Fruits	Strawberries 3.8g
	Bananas 1.4g
	Plums 2.3g
 Vegetables	Broccoli 2.8g
	Carrots 3.9g
	Sweet Potato 2.2g
 Legumes	Chickpeas 4.8g
	Lentils 3.8g
	Baked Beans 4.9g
 Nuts & Seeds	Almonds 15.9g
	Hazelnuts 9.4g
	Sunflower Seeds 7.1g
 Wholegrains	Oats 7.8g
	Wholegrain Bread 7g
	Wholewheat Pasta 4.4g

NUTRITION & HEALTH CLAIMS

Regulatory Landscape

Across the European Union (EU) and within the UK, foods can only carry a nutrition or health claim in accordance with EU* and UK** Regulations.

Foods can carry a 'source of fibre' or 'high fibre' nutrition claim if:

- Source of fibre - the product contains at least 3g of fibre per 100g or at least 1.5g of fibre per 100kcal.
- High fibre - the product contains at least 6g of fibre per 100g or at least 3g of fibre per 100kcal.

*Regulation (EC) No 1924/2006

**Nutrition (Amendment etc.) (EU Exit) Regulations 2020

All **health claims** on food products across the EU and within the UK require an assessment of scientific evidence by the European Food Safety Authority (EFSA) or the UK Nutrition and Health Claims Committee (UKNHCC), respectively.

Approved Health Claims

There are several approved health claims for dietary fibres relating to gastrointestinal health including⁸⁻¹¹:

Barley grain, oat grain, wheat bran and sugar beet fibres	Contributes to an increase in faecal bulk
Rye fibre	Contributes to normal bowel function
Wheat bran fibre	Contributes to an acceleration of intestinal transit time

FIBRE & HEALTH



Gastrointestinal Health

Consuming more dietary fibre has been shown to decrease intestinal transit time and exert faecal-bulking effects, soften stools and decrease constipation, in randomised-controlled trials.¹²



Colorectal Cancer

In prospective studies, high fibre intakes have been associated with a reduced risk of colorectal cancer, and the World Cancer Research Fund states there is strong 'probable' evidence that foods containing dietary fibre decreases the risk of colorectal cancer.^{13,14}



Other Non-Communicable Diseases

Further research suggests an association between increased fibre intake and reduced mortality risk and non-communicable diseases including cardiovascular disease, high blood pressure, type 2 diabetes, and some other cancers.¹⁵

REFERENCES

1. Codex Alimentarius Commission Joint FAO/WHO Food Standards Programme (2004) Proposals for a definition and methods of analysis for dietary fibre content.
2. EU (2011) Regulation (EU) No 1169/2011 of the European parliament and of the Council on the provision of food information to consumers. *Official Journal of the European Union* L304:18–63
3. PHE & FSA (2018) National Diet & Nutrition Survey: Results from Years 7 and 8 (combined) of the Rolling Programme (2014/2015 to 2015/2016).
4. IUN Alliance (2011) National Adult Nutrition Survey: Summary Report.
5. PHE (2016) Government Dietary Recommendations: Government recommendations for energy and nutrients for males and females aged 1–18 years and 19+ years.
6. FSAI (2011) Scientific Recommendations for Healthy Eating Guidelines in Ireland.
7. FSAI (2020) Scientific Recommendations for Food-Based Dietary Guidelines for 1 to 5 Year-Olds in Ireland.
8. EFSA NDA Panel (2011) *EFSA Journal* 9(6):2249
9. EFSA NDA Panel (2010) *EFSA Journal* 8(10):1817
10. EFSA NDA Panel (2011) *EFSA Journal* 9(12):2468
11. EFSA NDA Panel (2011) *EFSA Journal* 9(6):2258
12. SACN (2015) *Carbohydrates and Health*. London: TSO
13. Aune et al. (2011) *BMJ* 343: d6617
14. WCRF/AICR (2018) Continuous Update Project Expert Report 2018: Diet, Nutrition, Physical Activity and Colorectal Cancer.
15. Stephen et al. (2017) *Nutrition Research Reviews* 30: 149-190
16. De Filippo et al. (2010) *PNAS* 107(33) : 14691-14696
17. Koç et al. (2020) *Nutrition Bulletin* 45: 294-308

November 2020

FIND OUT MORE AT [YAKULT.CO.UK/HCP](https://www.yakult.co.uk/hcp)

Yakult is a science based company, dedicated to scientific research and education.

Contact us at science@yakult.co.uk or on **020 8842 7600** for more information.

This resource is intended for healthcare professionals and is not to be distributed to patients